

SHORT COMMUNICATION

First record of the sisal weevil, *Scyphophorus acupunctatus*, in Cyprus

VASSILIS VASSILIOU* AND PAVLOS KITSIS

Agricultural Research Institute, P.O. Box 22016, 1516, Nicosia, Cyprus

ABSTRACT

In May 2013, numerous adults of a coleopteran insect species that later was identified as the sisal weevil, *Scyphophorus acupunctatus* (Coleoptera: Curculionidae), were accidentally captured in both pitfall and funnel traps placed across Cyprus for the monitoring of red palm weevil, *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae). *S. acupunctatus* was found for the first time in the Germasogeia area of the Limassol district (34°71 81 N, 33°08 56 E) and in the Kissonerga area of the Paphos district (34°81 67 N, 32°40 00 E). During 2013 and early 2014, numerous adults of this species were also collected from red palm weevil traps from all over Cyprus.

KEY WORDS: Agavaceae, agave, dracaena, Dracenaceae, *Rhynchophorus ferrugineus*, trapping, yucca.

The sisal or agave weevil, *Scyphophorus acupunctatus* Gyllenhal (Coleoptera: Curculionidae) (syns: *S. anthraxcinus*, *S. interstitialis*, *S. robustior*, *Rhynchophorus asperulus*) (Coleoptera: Curculionidae), is a species native to Nearctic Region (USA, Mexico, Cayman islands, Costa Rica, Cuba, Netherlands Antilles, Dominican Republic, El Salvador, Guatemala, Haiti, Honduras, Jamaica, Nicaragua, Virgin islands, Belize, Brazil, Colombia, Venezuela). It has also been found in Africa (Kenya, South Africa, Tanzania), in Asia (Indonesia, Saudi Arabia) and Oceania (EPPO 2013). In Europe, it was first reported in 1980 in the Netherlands, on imported ornamental *Yucca* plants as well as several times on imported ornamental plant species such as the *Beaucarnea*, *Dasylyrion* and *Yucca* (van Rossem et al. 1981). It was

also recorded in Italy on *Beaucarnea recurvata* Lem. (Colombo 2000) and on *Agave americana* L. (EPPO 2008a), in France (EPPO 2008b), in Spain (Flinch and Zarazaga 2007) and in Greece (Kontodimas and Kallinikou 2010) on *A. americana* and *Agave* spp.

During 2013 and early 2014, sisal weevil was found in all the districts across Cyprus, in funnel and pitfall pheromone traps placed for the monitoring of red palm weevil *Rhynchophorus ferrugineus* Olivier (Coleoptera: Curculionidae) (Table 1). This insect species was identified at the Benaki Phytopathological Institute in Greece, using keys provided by Vaurie (1971). Apart from these captures, it was decided to conduct inspections in Cyprus in order to record any damages of hosts of this insect pest, mainly on the agaves and yuccas that are grown on the island.

*Corresponding author: email: vassilis@ari.gov.cy

TABLE 1. Number of adults of the sisal weevil, *Scyphophorus acupunctatus*, captured in red palm weevil pheromone traps across Cyprus during 2013 and early 2014.

Date of collection	Locality of collection	No of adults
May 2013	Limassol (Germasogeia area) (34°40 60 N, 33 4 60 E)	2
May 2013	Paphos (Kissonerga area) (34°49 30 N, 32 24 00 E)	8
May 2013	Paphos (Pegeia area) (34°88 33 N, 32 38 33 E)	2
June 2013	Larnaca (Zygi area) (34°73 33 N, 34 73 33 E)	5
June 2013	Larnaca (Mazotos village) (34°80 22 N, 33 49 00 E)	2
July 2013	Famagusta (Paralimni area) (35°03 33 N, 33 98 33 E)	5
October 2013	Nicosia (Kaimakli area) (35°18 91 N, 33 38 36 E)	1
January 2014	Limassol (Agios Tychonas area) (34°71 19 N, 33 12 69 E)	1
January 2014	Limassol (Germasogeia area) (34°40 60 N, 33 4 60 E)	2

According to the international literature, *S. acupunctatus* is a major pest of several ornamental species belonging to the Agavaceae and Dracenaceae families. It mainly attacks *Agave* plant species, including *A. sisalana* Perrine, *A. fourcroydes* Lem., *A. tequilana* F.A.C. Weber and *A. americana* L. (Vaurie 1971, Camino et al. 2002, Kontodimas and Kallinikou 2010, Setliff and Anderson 2011, EPPO 2013), the ponytail palm, *Beaucarnea recurvata* Lem., the Mexican grass tree, *Dasyllirion longissimum* Lem., the Canary islands dragon tree, *Dracaena draco* (L.), the giant Cabuya, *Furcraea foetida* (L.) Haw.

Additionally, this insect species is referred to attack various *Yucca* plant species such as the Spanish dagger, *Y. aloifolia* L., the spineless yucca, *Y. elephantipes* Lem., the soapweed yucca, *Y. glauca* Nutt., as well as the tuberose, *Polianthes tuberosa* L. (Vaurie 1971, Camino Lavin et al. 2002, Kontodimas and Kallinikou 2010, EPPO 2013) and the snake plant, *Sansevieria trifasciata* Prain (Pott 1976). This species is also considered a major problem in the tequila and henequen industries in Mexico and in the

sisal industries of Africa (Waring and Smith 1986).

According to the Department of Agriculture, Ministry of Agriculture, Natural Resources and Environment on Cyprus, the plant species mentioned in Table 2, are potential hosts of *S. acupunctatus* on Cyprus.

Adults of *S. acupunctatus* are small brown-black or black weevils without dorsal scales, 10-19mm long. They feed on leaves and bore into the bole of plant where females may oviposit from 25 to 30 eggs. Generally, there are 5 larval instars but their number may vary depending of the type of food that larvae were fed. The fully developed larva is about 18mm long, creamy white and legless. Pupation takes place within a cocoon made of plant fibres and debris. The total life cycle requires 50-90 days, with 4 or 5 generations per year (EPPO 2006). Male adults produce a pheromone that attracts both sexes (Montiel et al. 2009, Lopez-Martinez et al. 2011). Both larvae and adults are found in roots, lower leaves, and inside the heads (Montiel et al. 2008, CABI 2014).

TABLE 2. Potential host plants of sisal weevil, *Scyphophorus acupunctatus*, on Cyprus. Source: Department of Agriculture, Ministry of Agriculture, Natural Resources and Environment.

Species	Genus
<i>Beaucarnea recurvate</i> Lem.	Beaucarnea
<i>Yucca rostrata</i> Engelm. ex Trel.	Yucca
<i>Yucca gloriosa</i> L.	Yucca
<i>Agave americana</i> L.	Agave
<i>Agave americana</i> var. <i>variegata</i> Hook.	Agave
<i>Agave filifera</i> Salm-Dyck	Agave
<i>Agave attenuata</i> Salm-Dyck 1834	Agave
<i>Agave angustifolia</i> Haw.	Agave
<i>Agave parviflora</i> Torr.	Agave
<i>Dracaena draco</i> (L.) L.	Dracaena
<i>Dracaena marginata</i> Lam.	Dracaena
<i>Dracaena sanderiana</i> Sander ex Mast.	Dracaena
<i>Dracaena fragrans</i> (L.) Ker Gawl.	Dracaena

Adult damage consists of groups of feeding punctures on young leaves. In addition to feeding damage, the larvae may transfer bacteria that favour the development of secondary fungal or bacterial rots leading to premature death of the host. Most often, the infestation is not apparent until the damage is severe. Application of insecticides is the most commonly used suppression method for *S. acupunctatus* (Gonzalez et al. 2007).

To the best of our knowledge this is the first record of *Scyphophorus acupunctatus* from the island of Cyprus.

Acknowledgments

The authors are grateful to Dr Dimitris Kontodimas at the Benaki Phytopathological Institute in Athens, Greece, for

identifying this insect species. Also, the authors would like to thank Mr Efthymios Odysseos at the Department of Agriculture, Ministry of Agriculture, Natural Resources and Environment for his valuable contribution on the potential host plants of this insect species in Cyprus.

References

- CABI. 2014. Invasive Species Compendium. <http://www.cabi.org/isc/?compid=5&dsid=49421&loadmodule=datasheet&page=481&site=144>.
- Camino, L.M., G.V.R. Castrejon, B.R. Figueroa, L.L. Aldana and V.M.E. Estrada. 2002. *Scyphophorus acupunctatus* (Coleoptera: Curculionidae) attacking *Polianthes tuberosa*

- (Liliales: Agavaceae) in Morelos, Mexico. Fla. Entomol. 85: 392-393.
- Colombo, M. 2000. *Scyphophorus acupunctatus* Gyllenhal (Coleoptera: Curculionidae): prima segnalazione per l'Italia. Boll. Zool. Agrar. Bachic. 32: 165-170.
- EPPO. 2006. Deletions from the EPPO Alert List. http://www.eppo.int/QUARANTINE/Alert_List/deletions.htm?utm_source=www.eppo.org&utm_medium=int_redirect.
- EPPO. 2008a. *Scyphophorus acupunctatus* found in Sicilia, Italy. EPPO reporting service 2008/179.
- EPPO. 2008b. First report of *Scyphophorus acupunctatus* in France. EPPO reporting service 2008/220.
- EPPO. 2013. PQR database. Paris, France: European and Mediterranean Plant Protection Organization. <http://www.eppo.org>.
- Flinch, J.M. and M.A. Alonso-Zarazaga. 2007. El picudo negro de la pita o agave, o max del henequén, *Scyphophorus acupunctatus* Gyllenhal, 1838 (Coleoptera: Dryophthoridae): primera cita para la Península Ibérica. Bol. Soc. Entomol. Aragonesa. 41: 419-42.
- Gonzalez, H.H., J.I.L. del Real and J.F. Solis. 2007. Manejo de plagas del agave tequilero. Colegrio de Postgraduados, Montecillo, Mexico. 123 pp.
- Kontodimas, D.C. and E. Kallinikou. 2010. First record of the sisal weevil *Scyphophorus acupunctatus* (Coleoptera: Curculionidae) in Greece. Entomol. Hell. 19: 39-41.
- Lopez-Martinez, V., I.A. Tejacal, M. Atrande-Rodriguez, M.D.J. Garsia-Ramirez and J.C. Rojas. 2011. Daily activity of *Scyphophorus acupunctatus* (Coleoptera: Curculionidae) monitored with pheromone traps in a field of Mexican tuberose. Fla. Entomol. 94:1091-1093.
- Montiel, R.C., G.C. Guadalupe, J.C. Rojas, E.A. Malo, C.L. López, R. del Real and G.H. Hernández. 2008. Aggregation pheromone of the agave weevil, *Scyphophorus acupunctatus*. Entomol. Exp. Appl. 127: 207-217.
- Pott, J.N. 1976. A Yucca Borer, *Scyphophorus acupunctatus*, in Florida. Florida State Horticultural Society, pp. 414-416.
- Montiel, R.C., J.C. Rojas, L. Cruz-Lopez and H. Gonzalez-Hernandez. 2009. Factors affecting pheromone releases by *Scyphophorus acupunctatus* (Coleoptera: Curculionidae). Environ. Entomol. 38: 1423-1428.
- Setliff, G.P. and J.A. Anderson. 2011. First record of the agave weevil, *Scyphophorus acupunctatus* Gyllenhal (Coleoptera: Curculionidae: Dryophorinae), in Puerto Rico. Insecta Mundi. 0152: 1-3. <http://digitalcommons.unl.edu/cgi/viewcontent.cgi?article=1670&context=insectamundi>.
- Van Rossem, G., C.F. van de Bund, H.C. Burger and L.J.W. de Goffau. 1981. Bijzondere aantastingen door insecten in 1980. Entomol. Berichten. 41: 84-87.
- Vaurie, P. 1971. Review of *Scyphophorus* (Coleoptera: Rhynchophorinae). Coleopt. Bull. 25: 1-8.
- Waring, G.L. and R.I. Smith. 1986. Natural history and ecology of *Scyphophorus acupunctatus* (Coleoptera: Curculionidae) and its associated microbes in cultivated and native agaves. Ann. Entomol. Soc. Amer. 79: 334-340.

**μ Scyphophorus
acupunctatus**

*

, . . . 22016, 1516, ,

2013, μ

(pitfall), μ μ

μ

Rhynchophorus ferrugineus Olivier (Coleoptera:

Curculionidae)

, μ

Vaurie (1971),

Scyphophorus acupunctatus Gyllenhal (Coleoptera: Curculionidae). S.

acupunctatus

μ

μ

(34°71 81 N, 33 08 56 E)

(34°81 67 N, 32°40 00 E).

2013

μ

2014,

μ

μ

.